Nonlinear Systems

CS 205A:
Mathematical Methods for Robotics, Vision, and Graphics

Doug James (and Justin Solomon)

Part III: Nonlinear Problems

Not all numerical problems can be solved with \ in Matlab.

Question

Have we already seen a nonlinear problem?

Question

Have we already seen a nonlinear problem?

$$\begin{aligned} &\text{minimize } \|A\vec{x}\|_2 \\ &\text{such that } \|\vec{x}\|_2 = 1 \longleftarrow \text{nonlinear!} \end{aligned}$$

Root-Finding Problem

$$\text{Given: } f : \mathbb{R}^n \to \mathbb{R}^m$$

$$\text{Find: } \vec{x}^* \text{ with } f(\vec{x}^*) = \vec{0}$$

Issue: Regularizing Assumptions

$$f(x) = \begin{cases} -1 & \text{when } x \leq 0 \\ 1 & \text{when } x > 0 \end{cases}$$

Issue: Regularizing Assumptions

$$f(x) = \begin{cases} -1 & \text{when } x \leq 0 \\ 1 & \text{when } x > 0 \end{cases}$$

$$g(x) = \begin{cases} -1 & \text{when } x \in \mathbb{Q} \\ 1 & \text{when } x \notin \mathbb{Q} \end{cases}$$

Typical Regularizing Assumptions

Continuous

$f(\vec{x}) \to f(\vec{y})$ as $\vec{x} \to \vec{y}$

Typical Regularizing Assumptions

Continuous

$f(\vec{x}) \to f(\vec{y})$ as $\vec{x} \to \vec{y}$

Lipschitz

$\|f(\vec{x}) - f(\vec{y})\|_2 \leq c\|\vec{x} - \vec{y}\|_2$ for all $\vec{x}, \vec{y}$ (same c)

Typical Regularizing Assumptions

Continuous

$f(\vec{x}) \to f(\vec{y})$ as $\vec{x} \to \vec{y}$

Lipschitz

$\|f(\vec{x}) - f(\vec{y})\|_2 \leq c\|\vec{x} - \vec{y}\|_2$ for all $\vec{x}, \vec{y}$ (same c)

Differentiable

$Df(\vec{x})$ exists for all $\vec{x}$

Typical Regularizing Assumptions

Continuous

$f(\vec{x}) \to f(\vec{y})$ as $\vec{x} \to \vec{y}$

Lipschitz

$\|f(\vec{x}) - f(\vec{y})\|_2 \leq c\|\vec{x} - \vec{y}\|_2$ for all $\vec{x}, \vec{y}$ (same c)

Differentiable

$Df(\vec{x})$ exists for all $\vec{x}$

C^k

k derivatives exist and are continuous

Today

$$f : \mathbb{R} \to \mathbb{R}$$

Property of Continuous Functions

Intermediate Value Theorem

Suppose that $f : [a, b] \to \mathbb{R}$ is continuous and that $f(a) < u < f(b)$ or $f(b) < u < f(a)$. Then, there exists $z \in (a, b)$ such that $f(z) = u$

Reasonable Input

- Continuous function $f(x)$
- $\ell, r \in \mathbb{R}$ with $f(\ell) \cdot f(r) < 0$ (why?)

Bisection Algorithm

1. Compute $c = {}^{\ell+r}/_{2}$.
2. If $f(c) = 0$, return $x^* = c$.
3. If $f(\ell) \cdot f(c) < 0$, take $r \leftarrow c$. Otherwise take $\ell \leftarrow c$.
4. Return to step 1 until $|r - \ell| < \varepsilon$; then return c.

Bisection: Illustration



Two Important Questions

1. Does it converge?

Two Important Questions

1. Does it converge?

   Yes! Unconditionally.

Two Important Questions

1. Does it converge?
   Yes! Unconditionally.

2. How quickly?

Convergence Analysis

Examine E_k with

$$|x_k - x^*| < E_k.$$

Bisection: Linear Convergence

$$E_{k+1} \leq \frac{1}{2} E_k$$

for $E_k \equiv |r_k - \ell_k|$

Fixed Points

$$g(x^*) = x^*$$

Fixed Points

$$g(x^*) = x^*$$

Question:

Same as root-finding?

Simple Strategy

$$x_{k+1} = g(x_k)$$

Convergence Criterion

$$
\begin{aligned}
E_k &\equiv |x_k - x^*| \\
&= |g(x_{k-1}) - g(x^*)|
\end{aligned}
$$

Convergence Criterion

$$
\begin{aligned}
E_k &\equiv |x_k - x^*| \\
&= |g(x_{k-1}) - g(x^*)| \\
&\leq c|x_{k-1} - x^*| \\
&\qquad\qquad \text{if } g \text{ is Lipschitz} \\
&= cE_{k-1}
\end{aligned}
$$

Convergence Criterion

$$
\begin{aligned}
E_k &\equiv |x_k - x^*| \\
&= |g(x_{k-1}) - g(x^*)| \\
&\leq c|x_{k-1} - x^*| \\
&\qquad\qquad \text{if } g \text{ is Lipschitz} \\
&= cE_{k-1} \\
\implies E_k &\leq c^k E_0 \\
&\to 0 \text{ as } k \to \infty \quad (c < 1)
\end{aligned}
$$

Alternative Criterion

Lipschitz *near* x^* with good starting point.

Alternative Criterion

Lipschitz *near* x^* with good starting point.

e.g. C^1 with $|g'(x^*)| < 1$

Convergence Rate of Fixed Point

When it converges...

Always linear (why?)

Convergence Rate of Fixed Point

When it converges...
Always linear (why?)

Often quadratic! ($\rightarrow$ board)

Approach for Differentiable $f(x)$



Newton's Method

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

Newton's Method

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

Fixed point iteration on

$$g(x) \equiv x - \frac{f(x)}{f'(x)}$$

Convergence of Newton

Simple Root

A root x^* with $f'(x^*) \neq 0$.

Convergence of Newton

Simple Root

A root x^* with $f'(x^*) \neq 0$.

Quadratic convergence in this case! ($\rightarrow$ board)

Issue

Differentiation is hard!

Secant Method

$$x_{k+1} = x_k - \frac{f(x_k)(x_k - x_{k-1})}{f(x_k) - f(x_{k-1})}$$

Secant Method

$$x_{k+1} = x_k - \frac{f(x_k)(x_k - x_{k-1})}{f(x_k) - f(x_{k-1})}$$

Trivia:

Converges at rate $\frac{1+\sqrt{5}}{2} \approx 1.6180339887\ldots$

Secant Method

$$x_{k+1} = x_k - \frac{f(x_k)(x_k - x_{k-1})}{f(x_k) - f(x_{k-1})}$$

Trivia:

Converges at rate $\frac{1+\sqrt{5}}{2} \approx 1.6180339887\ldots$

("Golden Ratio")

Hybrid Methods

Want: Convergence rate of secant/Newton with convergence guarantees of bisection

Hybrid Methods

Want: Convergence rate of secant/Newton with convergence guarantees of bisection

e.g. **Dekker's Method:** Take secant step if it is in the bracket, bisection step otherwise

Single-Variable Conclusion

- Unlikely to solve exactly, so we settle for iterative methods
- Must check that method converges at all
- Convergence rates:
  - Linear: $E_{k+1} \leq CE_k$ for some $0 \leq C < 1$
  - Superlinear: $E_{k+1} \leq CE_k^r$ for some $r > 1$
  - Quadratic: $r = 2$
  - Cubic: $r = 3$
- Time *per* iteration also important

Next