

Radiosity

Classic radiosity = finite element method

Assumptions

- Diffuse reflectance
- Usually polygonal surfaces

Advantages

- View independent solution

Techniques

- Meshing
- Form factors
- Solving linear equations

CS348B Lecture 17

Pat Hanrahan, Spring 2001

First Radiosity Pictures ...



Parry Moon and Domina Spencer (MIT), Lighting Design, 1948

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Early Radiosity




CS348B Lecture 17

Pat Hanrahan, Spring 2001

The Radiosity Equation

Assume diffuse reflection only

Solve for radiosity (2D function)



$$B(x) = B_e(x) + \rho(x)E(x)$$


$$B(x) = B_e(x) + \rho(x) \int_{M^2} F(x, x') B(x') dA'$$

$$F(x, x') = \frac{G(x, x')}{\pi} = \frac{\cos \theta \cos \theta'}{\pi \|x - x'\|^2} V(x, x')$$

CS348B Lecture 17

Pat Hanrahan, Spring 2001


Classic Radiosity Algorithm



CS348B Lecture 17

Pat Hanrahan, Spring 2001

Simple Room Scene



Example from John Wallace

CS348B Lecture 17

Pat Hanrahan, Spring 2001


Derivation

Radiosity integral equation

$$B(x) = B_e(x) + \rho(x) \int_{M^2} B(x') F(x, x') dA'$$

Piecewise constant basis functions

$$B(x) = \sum_i B_i N_i(x)$$



$$B_e(x) = \sum_i E_i N_i(x)$$



$$\rho(x) = \sum_i \rho_i N_i(x)$$



CS348B Lecture 17

Pat Hanrahan, Spring 2001

Derivation

Convert integral equation to matrix equation

$$B(x) = B_e(x) + \rho(x) \int_{M^2} F(x, x') \sum_j B_j N_j(x') dA'$$

$$\sum_i B_i N_i(x) = \sum_i B_i N_i(x) + \sum_i \rho_i N_i(x) \left[\sum_j B_j \int_{M^2} F(x, x') N_j(x') dA' \right]$$

$$\int \left(\sum_i B_i N_i(x) = \sum_i B_i N_i(x) + \sum_i \rho_i N_i(x) \left[\sum_j B_j \int_{M^2} F(x, x') N_j(x') dA' \right] \right) dA$$

$$B_i A_i = E_i A_i + \rho_i \sum_j B_j \int_{M^2} \int_{M^2} F(x, x') N_i(x) N_j(x') dA dA'$$

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Form Factor

Throughput

$$T_{ij} = T_{ji} = \int \int_{A_i A_j} \frac{\cos \theta'_o \cos \theta_i}{\pi \|x - x'\|^2} V(x, x') dA dA'$$

Reciprocity

$$\begin{aligned} T_{ij} &\equiv A_i F_{ij} \\ T_{ji} &\equiv A_j F_{ji} \end{aligned} \quad \Rightarrow \quad A_i F_{ij} = A_j F_{ji}$$

Summation

$$\sum_j F_{ij} = \sum_i F_{ji} = 1$$

Form factor is the percentage of light ...

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Classic Radiosity

Power Balance

$$B_i A_i = E_i A_i + \rho_i \sum_j B_j A_j F_{ji}$$

Reciprocity

$$A_i F_{ij} = A_j F_{ji} \Rightarrow B_i = E_i + \rho_i \sum_j F_{ij} B_j$$

Radiosity System

$$\begin{pmatrix} 1 - \rho_1 F_{11} & -\rho_1 F_{12} & \cdots & -\rho_1 F_{1n} \\ -\rho_2 F_{21} & 1 - \rho_2 F_{22} & \cdots & -\rho_2 F_{21} \\ \vdots & \vdots & \ddots & \vdots \\ -\rho_n F_{n1} & -\rho_n F_{n2} & \cdots & 1 - \rho_n F_{nn} \end{pmatrix} \begin{pmatrix} B_1 \\ B_2 \\ \vdots \\ B_n \end{pmatrix} = \begin{pmatrix} E_1 \\ E_2 \\ \vdots \\ E_n \end{pmatrix}$$

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Form Factors

Differential-differential - $F(x, x') dA'$

$$F_{dA_i, dA_j} = \frac{\cos \theta'_o \cos \theta_i}{\pi \|x - x'\|^2} V(x, x') dA'$$

Differential-finite - $E_j(x)$

$$F_{dA_i, A_j} = \int_{A_j} \frac{\cos \theta'_o \cos \theta_i}{\pi \|x - x'\|^2} V(x, x') dA'$$


Finite-finite - $\Phi_{i,j}$

$$F_{A_i, A_j} = \frac{1}{A_i} \int_{A_i} \int_{A_j} \frac{\cos \theta'_o \cos \theta_i}{\pi \|x - x'\|^2} V(x, x') dA dA'$$

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Analytical Form Factors



$$X = \frac{a}{c}$$

$$Y = \frac{b}{c}$$

$$F_{A_1, A_2} = \frac{2}{\pi X Y} \left\{ \ln \left[\frac{(1+X^2)(1+Y^2)}{(1+X^2+Y^2)} \right]^{1/2} + X \sqrt{1+Y^2} \tan^{-1} \frac{X}{\sqrt{1+Y^2}} \right. \\ \left. + Y \sqrt{1+X^2} \tan^{-1} \frac{Y}{\sqrt{1+X^2}} - X \tan^{-1} X - Y \tan^{-1} Y \right\}$$



CS348B Lecture 17

Pat Hanrahan, Spring 2001

Hemicube Algorithm

First radiosity algorithm to deal with occlusion

1. Render scene from the point of view of each vertex/element
2. Compute delta form factors – contribution from each pixel




Typical resolution: 32x32

CS348B Lecture 17


Pat Hanrahan, Spring 2001

Hemicube Delta Form Factors



$$\cos \phi = \frac{1}{\sqrt{x^2 + y^2 + 1}}$$

$$\Delta F = \frac{\Delta A}{\pi(x^2 + y^2 + 1)^2}$$



$$\cos \phi = \frac{1}{\sqrt{1 + y^2 + z^2}}$$

$$\Delta F = \frac{\Delta A}{\pi(1 + y^2 + z^2)^2}$$

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Hemicube Algorithms

Advantages

- + First practical method -> Patent!
- + Use existing rendering systems; Hardware!
- + Computes all form factors in $O(n)$

Disadvantages

- Computes differential-finite form factor
- Aliasing errors due to sampling
Randomly rotate/shear hemicube
- Proximity errors
- Visibility errors
- Expensive to compute a single form factor

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Solve $[F][B] = [E]$

Direct methods: $O(n^3)$

- Gaussian elimination

Goral, Torrance, Greenberg, Battaile, 1984

Iterative methods: $O(n^2)$

Convergence

Energy conservation -> diagonally dominant -> converge

- Gauss-Seidel, Jacobi: Gathering

Nishita, Nakamae, 1985

Cohen, Greenberg, 1985

- Southwell: Shooting

Cohen, Chen, Wallace, Greenberg, 1988

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Iterative Solvers

Iteration


$$(I - F)^{-1}B = E$$
$$B = (I + B + B^2 + \dots)E$$

$$B^0 = E$$
$$B^1 = E + FB^0$$
$$\dots$$
$$B^n = E + FB^{n-1}$$

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Gathering




```
for(i=0; i<n; i++)  
    B[i] = Be[i];  
  
while( !converged ) {  
    for(i=0; i<n; i++) {  
        E[i] = 0;  
        for(j=0; j<n; j++)  
            E[i] += F[i][j] * B[j];  
        B[i] = rho[i]*E[i];  
    }  
}
```

Row of F times B
Calculate one row of F and discard

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Shooting



Brightness order


Column of F times B

```
for(i=0; i<n; i++)  
    B[i] = dB[i] = Be[i];  
while( !converged ) {  
    set i st dB[i] is the largest;  
    for(j=0;j<n;j++)  
        if(i!=j) {  
            dBj = rho[j]*F[j][i]*dB[i];  
            dB[j] += dBj;  
            B[j] += dBj;  
        }  
    dB[i]=0;  
}
```

CS348B Lecture 17

Pat Hanrahan, Spring 2001

Results: Gathering vs. Shooting




From Cohen et al.

Figure 5.9: Convergence versus number of steps for three algorithms.


CS348B Lecture 17

Pat Hanrahan, Spring 2001

Accuracy



Reference Solution




Uniform Mesh

Table in room sequence from Cohen and Wallace


CS348B Lecture 17

Pat Hanrahan, Spring 2001

Artifacts



- A. Blocky shadows
- B. Missing features
- C. Mach bands
- D. Inappropriate shading discontinuities
- E. Unresolved discontinuities






Error Image

CS348B Lecture 17

Pat Hanrahan, Spring 2001



Increasing Resolution



CS348B Lecture 17

Pat Hanrahan, Spring 2001


Adaptive Meshing



CS348B Lecture 17

Pat Hanrahan, Spring 2001

Megaradiosity



From S. Teller, T. Funkhouser, C. Fowler, P. Hanrahan

CS348B Lecture 17

Pat Hanrahan, Spring 2001