

Reflection Models I

Today

- Types of reflection models
- The BRDF and reflectance
- The reflection equation
- Ideal reflection and refraction
- Fresnel effect
- Ideal diffuse

Next lecture

- Glossy and specular reflection models
- Rough surfaces and microfacets, self-shadowing
- Anisotropic reflection models

CS348B Lecture 10

Pat Hanrahan, Spring 2006

Reflection Models

Definition: Reflection is the process by which light incident on a surface interacts with the surface such that it leaves on the incident side without change in frequency.

Properties

- Spectra and Color [Moon Spectra]
- Polarization
- Directional distribution


CS348B Lecture 10

Pat Hanrahan, Spring 2006

Types of Reflection Functions


Ideal Specular

- Reflection Law
- Mirror




Ideal Diffuse

- Lambert's Law
- Matte



Specular

- Glossy
- Directional diffuse



CS348B Lecture 10

Pat Hanrahan, Spring 2006


Materials



Plastic



Metal




Matte

From Apodaca and Gritz, *Advanced RenderMan*

CS348B Lecture 10

Pat Hanrahan, Spring 2006


Spheres [Matusik et al.]



CS348B Lecture 10

Pat Hanrahan, Spring 2006

The Reflection Equation




$$L_r(x, \omega_r) = \int_{H^2} f_r(x, \omega_i \rightarrow \omega_r) L_i(x, \omega_i) \cos \theta_i d\omega_i$$

CS348B Lecture 10

Pat Hanrahan, Spring 2006

The BRDF

Bidirectional Reflectance-Distribution Function




$$f_r(\omega_i \rightarrow \omega_r) \equiv \frac{dL_r(\omega_i \rightarrow \omega_r)}{dE_i} \left[\frac{1}{sr} \right]$$

CS348B Lecture 10

Pat Hanrahan, Spring 2006

The BSSRDF

Bidirectional Surface Scattering Reflectance-Distribution Function





$$S(x_i, \omega_i \rightarrow x_r, \omega_r) \equiv \frac{dL_r(x_i, \omega_i \rightarrow x_r, \omega_r)}{d\Phi_i}$$

CS348B Lecture 10

Pat Hanrahan, Spring 2006

Gonioreflectometer




CS348B Lecture 10

Pat Hanrahan, Spring 2006

Properties of BRDF's


1. Linear



From Sillion, Arvo, Westin, Greenberg

2. Reciprocity principle

$$f_r(\omega_r \rightarrow \omega_i) = f_r(\omega_i \rightarrow \omega_r)$$




CS348B Lecture 10

Pat Hanrahan, Spring 2006

Properties of BRDF's

3. Isotropic vs. anisotropic

$$f_r(\theta_i, \varphi_i; \theta_r, \varphi_r) = f_r(\theta_i, \theta_r, \varphi_r - \varphi_i)$$



Reciprocity and isotropy

$$f_r(\theta_i, \theta_r, \varphi_r - \varphi_i) = f_r(\theta_r, \theta_i, \varphi_i - \varphi_r) = f_r(\theta_i, \theta_r, |\varphi_r - \varphi_i|)$$


4. Energy conservation

CS348B Lecture 10

Pat Hanrahan, Spring 2006

Energy Conservation

$$\begin{aligned} \frac{d\Phi_r}{d\Phi_i} &= \frac{\int_{\Omega_r} L_r(\omega_r) \cos \theta_r d\omega_r}{\int_{\Omega_i} L_i(\omega_i) \cos \theta_i d\omega_i} \\ &= \frac{\int_{\Omega_r} \int_{\Omega_i} f_r(\omega_i \rightarrow \omega_r) L_i(\omega_i) \cos \theta_i d\omega_i \cos \theta_r d\omega_r}{\int_{\Omega_i} L_i(\omega_i) \cos \theta_i d\omega_i} \\ &\leq 1 \end{aligned}$$




CS348B Lecture 10

Pat Hanrahan, Spring 2006

The Reflectance

Definition: Reflectance is ratio of reflected to incident power



$$\rho(\Omega_i \rightarrow \Omega_r) \equiv \frac{\int \int f_r(\omega_i \rightarrow \omega_r) \cos \theta_i d\omega_i \cos \theta_r d\omega_r}{\int \cos \theta_i d\omega_i}$$
$$\leq \frac{\int \int f_r(\omega_i \rightarrow \omega_r) L_i(\omega_i) \cos \theta_i d\omega_i \cos \theta_r d\omega_r}{\int L_i(\omega_i) \cos \theta_i d\omega_i}$$

Conservation of energy: $0 < \rho < 1$


3 by 3 set of possibilities: $\{d\omega_i, \Omega_i, H_i^2\} \times \{d\omega_r, \Omega_r, H_r^2\}$

Units: ρ [dimensionless], f_r [1/steradians]

CS348B Lecture 10

Pat Hanrahan, Spring 2006

Law of Reflection




$$\hat{R} + (-\hat{I}) = 2 \cos \theta \hat{N} = -2(\hat{I} \bullet \hat{N}) \hat{N}$$

$$\hat{R} = \hat{I} - 2(\hat{I} \bullet \hat{N}) \hat{N}$$

CS348B Lecture 10

Pat Hanrahan, Spring 2006

Ideal Reflection (Mirror)



$$L_{r,m}(\theta_r, \varphi_r) = L_i(\theta_r, \varphi_r \pm \pi)$$



$$f_{r,m}(\theta_i, \varphi_i; \theta_r, \varphi_r) = \frac{\delta(\cos \theta_i - \cos \theta_r)}{\cos \theta_i} \delta(\varphi_i - \varphi_r \pm \pi)$$

$$\begin{aligned} L_{r,m}(\theta_r, \varphi_r) &= \int f_{r,m}(\theta_i, \varphi_i; \theta_r, \varphi_r) L_i(\theta_i, \varphi_i) \cos \theta_i d\cos \theta_i d\varphi_i \\ &= \int \frac{\delta(\cos \theta_i - \cos \theta_r)}{\cos \theta_i} \delta(\varphi_i - \varphi_r \pm \pi) L_i(\theta_i, \varphi_i) \cos \theta_i d\cos \theta_i d\varphi_i \\ &= L_i(\theta_r, \varphi_r \pm \pi) \end{aligned}$$

CS348B Lecture 10

Pat Hanrahan, Spring 2006

Snell's Law



$$\varphi_t = \varphi_i \pm \pi$$


$$n_i \sin \theta_i = n_t \sin \theta_t$$

$$n_i \hat{\mathbf{N}} \times \hat{\mathbf{I}} = n_t \hat{\mathbf{N}} \times \hat{\mathbf{T}}$$

CS348B Lecture 10

Pat Hanrahan, Spring 2006

Law of Refraction



$$\hat{N} \times \hat{T} = \mu \hat{N} \times \hat{I}$$

$$\hat{N} \times (\hat{T} - \mu \hat{I}) = 0$$

$$\hat{T} = \mu \hat{I} + \gamma \hat{N}$$

$$\hat{T}^2 = 1 = \mu^2 + \gamma^2 + 2\mu\gamma \hat{I} \bullet \hat{N}$$

$$\begin{aligned}\gamma &= -\mu \hat{I} \bullet \hat{N} \pm \left\{ 1 - \mu^2 \left(1 - (\hat{I} \bullet \hat{N})^2 \right) \right\}^{1/2} \\ &= \mu \cos \theta_i \pm \left\{ 1 - \mu^2 \sin^2 \theta_i \right\}^{1/2}\end{aligned}$$

Total internal reflection:

$$1 - \mu^2 (1 - (\hat{I} \bullet \hat{N})^2) < 0$$


$$\begin{aligned}&= \mu \cos \theta_i \pm \cos \theta_t \quad \leftarrow \gamma = \mu - 1 \\ &= \mu \cos \theta_i - \cos \theta_t\end{aligned}$$

CS348B Lecture 10

Pat Hanrahan, Spring 2006

Optical Manhole

Total internal reflection



$$n_w = \frac{4}{3}$$




From Livingston and Lynch

CS348B Lecture 10

Pat Hanrahan, Spring 2006


Fresnel Reflectance

Metal (Aluminum)



Gold $F(0)=0.82$
Silver $F(0)=0.95$

Dielectric ($N=1.5$)



Glass $n=1.5 F(0)=0.04$
Diamond $n=2.4 F(0)=0.15$

Schlick Approximation $F(\theta) = F(0) + (1 - F(0))(1 - \cos \theta)^5$

CS348B Lecture 10

Pat Hanrahan, Spring 2006

Experiment

Reflections from a shiny floor



From Lafontaine, Foo, Torrance, Greenberg, SIGGRAPH 97



CS348B Lecture 10

Pat Hanrahan, Spring 2006

Cook-Torrance Model for Metals

Reflectance of Copper as a function of wavelength and angle of incidence

Light spectra



Measured Reflectance



Approximated Reflectance

Cook-Torrance approximation


$$R = R(0) + R(\pi/2) \left[\frac{F(\theta) - F(0)}{F(\pi/2) - F(0)} \right]$$

CS348B Lecture 10

Pat Hanrahan, Spring 2006

Ideal Diffuse Reflection

Assume light is equally likely to be reflected in any output direction



$$\begin{aligned} L_{r,d}(\omega_r) &= \int f_{r,d} L_i(\omega_i) \cos \theta_i d\omega_i \\ &= f_{r,d} \int L_i(\omega_i) \cos \theta_i d\omega_i \\ &= f_{r,d} E \end{aligned}$$

$$M = \int L_r(\omega_r) \cos \theta_r d\omega_r = L_r \int \cos \theta_r d\omega_r = \pi L_r$$

$$\rho_d = \frac{M}{E} = \frac{\pi L_r}{E} = \frac{\pi f_{r,d} E}{E} = \pi f_{r,d} \Rightarrow f_{r,d} = \frac{\rho_d}{\pi}$$

Lambert's Cosine Law $M = \rho_d E = \rho_d E_s \cos \theta_s$

CS348B Lecture 10

Pat Hanrahan, Spring 2006

“Diffuse” Reflection

Theoretical

- **Bouguer - Special micro-facet distribution**
- **Seeliger - Subsurface reflection**
- **Multiple surface or subsurface reflections**

Experimental

- **Pressed magnesium oxide powder**
- **Almost never valid at high angles of incidence**

Paint manufacturers attempt to create ideal diffuse