

The Rendering Equation

Direct (*local*) illumination

- Light directly from light sources
- No shadows

Indirect (*global*) illumination

- Hard and soft shadows
- Diffuse interreflections (radiosity)
- Glossy interreflections (caustics)

CS348B Lecture 13

Pat Hanrahan, Spring 2006




Radiosity



CS348B Lecture 13

Pat Hanrahan, Spring 2006

Lighting Effects




CS348B Lecture 13

Pat Hanrahan, Spring 2006

Challenge

**To evaluate the reflection equation
the incoming radiance must be known**

$$L_r(x, \omega_r) = \int_{H^2} f_r(x, \omega_i \rightarrow \omega_r) L_i(x, \omega_i) \cos \theta_i d\omega_i$$



**To evaluate the incoming radiance
the reflected radiance must be known**

CS348B Lecture 13

Pat Hanrahan, Spring 2006

To The Rendering Equation


Questions

- 1. How is light measured?**
- 2. How is the spatial distribution of light energy described?**
- 3. How is reflection from a surface characterized?**
- 4. What are the conditions for equilibrium flow of light in an environment?**

CS348B Lecture 13

Pat Hanrahan, Spring 2006

The Grand Scheme



CS348B Lecture 13

Pat Hanrahan, Spring 2006

Balance Equation

Accountability

$$[\text{outgoing}] - [\text{incoming}] = [\text{emitted}] - [\text{absorbed}]$$

■ Macro level

The total light energy put into the system must equal the energy leaving the system (usually, via heat).

$$\Phi_o - \Phi_i = \Phi_e - \Phi_a$$

■ Micro level

The energy flowing into a small region of phase space must equal the energy flowing out.

$$B(x) - E(x) = B_e(x) - E_a(x)$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Surface Balance Equation


$$[\text{outgoing}] = [\text{emitted}] + [\text{reflected}]$$

$$\begin{aligned} L_o(x, \omega_o) &= L_e(x, \omega_o) + L_r(x, \omega_o) \\ &= L_e(x, \omega_o) + \int_{H^2} f_r(x, \omega_i \rightarrow \omega_o) L_i(x, \omega_i) \cos \theta_i d\omega_i \end{aligned}$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Direction Conventions




CS348B Lecture 13

Pat Hanrahan, Spring 2006

Surface Balance Equation

[outgoing] = [emitted] + [reflected] + [transmitted]



CS348B Lecture 13

Pat Hanrahan, Spring 2006


Two-Point Geometry

$$\omega(x, x') = \omega(x \rightarrow x') = \frac{x' - x}{|x' - x|}$$

Ray Tracing

$$x^*(x, \omega)$$

$$\omega_i = \omega(x, x')$$




CS348B Lecture 13

Pat Hanrahan, Spring 2006

Coupling Equations

$$L(x', \omega') = L_o(x', \omega_o)$$



Invariance of radiance

$$L_i(x, \omega_i) = L(x, -\omega_i)$$

$$L(x, \omega) = L(x', \omega')$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

The Rendering Equation

Directional form

$$L(x, \omega) = L_e(x, \omega) + \int f_r(x, \omega' \rightarrow \omega) L(x^*(x, \omega'), -\omega') \cos \theta' d\omega'$$

Integrate over hemisphere of directions **Transport operator i.e. ray tracing**

CS348B Lecture 13

Pat Hanrahan, Spring 2006

The Rendering Equation


Surface form

$$L(x', x) = L_e(x', x) + \int f_r(x'', x', x) L(x'', x') G(x'', x') dA''(x'')$$

Integrate over all surfaces **Geometry term**

$$G(x'', x') = \frac{\cos \theta_i'' \cos \theta_o'}{\|x'' - x'\|^2} V(x'', x')$$

Visibility term

$$V(x'', x') = \begin{cases} 1 & \text{visible} \\ 0 & \text{not visible} \end{cases}$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

The Radiosity Equation

Assume diffuse reflection

1. $f_r(x, \omega_i \rightarrow \omega_o) = f_r(x) \Rightarrow \rho(x) = \pi f_r(x)$

2. $L(x, \omega) = B(x) / \pi$

$$B(x) = B_e(x) + \rho(x)E(x)$$

$$B(x) = B_e(x) + \rho(x) \int F(x, x') B(x') dA'(x')$$



$$F(x, x') = \frac{G(x, x')}{\pi}$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Integral Equations

Integral equations of the 1st kind

$$f(x) = \int k(x, x') g(x') dx'$$

Integral equations of the 2nd kind

$$f(x) = g(x) + \int k(x, x') f(x') dx'$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Linear Operators

Linear operators act on functions like matrices act on vectors

$$h(x) = (L \circ f)(x)$$

They are linear in that

$$L \circ (af + bg) = a(L \circ f) + b(L \circ g)$$

Types of linear operators

$$(K \circ f)(x) \equiv \int k(x, x') f(x') dx'$$

$$(D \circ f)(x) \equiv \frac{\partial f}{\partial x}(x)$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Solving the Rendering Equation

Rendering Equation

$$L = L_e + K \circ L$$

$$(I - K) \circ L = L_e$$

Solution

$$L = (I - K)^{-1} \circ L_e$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Formal Solution

Neumann series

$$(I - K)^{-1} = \frac{1}{I - K} = I + K + K^2 + \dots$$

Verify

$$\begin{aligned}(I - K) \circ (I - K)^{-1} &= (I - K) \circ (I + K + K^2 + \dots) \\ &= (I + K + \dots) - (K + K^2 + \dots) \\ &= I\end{aligned}$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Successive Approximations

Successive approximations

$$L^1 = L_e$$

$$L^2 = L_e + K \circ L^1$$

...

$$L^n = L_e + K \circ L^{n-1}$$


Converged

$$L^n = L^{n-1} \quad \therefore \quad L^n = L_e + K \circ L^n$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Successive Approximation




CS348B Lecture 13

Pat Hanrahan, Spring 2006

Light Path

$$S(x_0, x_1) = L_e(x_0, x_1)$$




$$L_S(x_0, x_1, x_2, x_3)$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Light Path




$$L_s(x_0, x_1, x_2, x_3) = S(x_0, x_1)G(x_1, x_2)f_r(x_0, x_1, x_2)G(x_1, x_2)f_r(x_1, x_2, x_3)$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Light Paths



$$L(x_2, x_3) = \int_{A_0} \int_{A_1} L_s(x_0, x_1, x_2, x_3) dA(x_0) dA(x_1)$$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Light Transport

Integrate over all paths of all lengths

$$L(x_{k-1}, x_k)$$

$$= \sum_{k=1}^{\infty} \int_{M^2} \cdots \int_{M^2} L_S(x_0, \dots, x_{k-2}, x_{k-1}, x_k) dA(x_0) \cdots dA(x_{k-2})$$


Question:

- How to sample space of paths?

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Classic Ray Tracing




Forward (from eye): $E S^* (D|G) L$

CS348B Lecture 13

Pat Hanrahan, Spring 2006

Photon Paths



CS348B Lecture 13

Pat Hanrahan, Spring 2006

How to Solve It?

Finite element methods

- **Classic radiosity**
 - Mesh surfaces
 - Piecewise constant basis functions
 - Solve matrix equation
- **Not practical for rendering equation**

Monte Carlo methods

- **Path tracing (distributed ray tracing)**
- **Bidirectional ray tracing**
- **Photon mapping**

CS348B Lecture 13

Pat Hanrahan, Spring 2006